



**EDGEWOOD**

RESEARCH DEVELOPMENT & ENGINEERING CENTER

U.S. ARMY CHEMICAL AND BIOLOGICAL DEFENSE COMMAND

AD-A285 920



ERDEC-TR-202

**AQUATIC TOXICITY  
OF DECONTAMINATING SOLUTIONS DS-2/DS-2P**

DTIC  
ELECTE  
NOV 04 1994  
S B D

M.V. Haley  
N.A. Chester  
C.W. Kurnas  
C.T. Phillips

RESEARCH AND TECHNOLOGY DIRECTORATE

94-34307



1788

September 1994

Approved for public release; distribution is unlimited.



Aberdeen Proving Ground, MD 21010-5423

94 11 3 05 9

**Disclaimer**

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 1994 September	3. REPORT TYPE AND DATES COVERED Final, 93 May - 93 Dec		
4. TITLE AND SUBTITLE Aquatic Toxicity of Decontaminating Solutions DS-2/DS-2P		5. FUNDING NUMBERS PR-10464806DF97 Sales Order No. 2FK4		
6. AUTHOR(S) Haley, M.V.; Chester, N.A.; Kurnas, C.W.; and Phillips, C.T.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) DIR, ERDEC, ATTN: SCBRD-RTL, APG, MD 21010-5423		8. PERFORMING ORGANIZATION REPORT NUMBER ERDEC-TR-202		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words)  The U.S. Army is working on reducing the possibility of human health effects from exposure to the decontaminating solution DS-2. One of the components of the DS-2 mixture, ethylene glycol monomethyl ether (EGME), has been determined to cause birth defects, fetotoxicity, and bone marrow problems in laboratory animals. The proposed formulation (DS-2P) replaces EGME with propylene glycol monomethyl ether. This report provides initial environmental screening results using <i>Daphnia magna</i> , fathead minnow, <i>Photobacterium phosphoreum</i> , and earthworm.				
14. SUBJECT TERMS Aquatic toxicology <i>Daphnia magna</i> Microtox		Earthworm Fathead minnow Decon solution		15. NUMBER OF PAGES 16
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		16. PRICE CODE
19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED		20. LIMITATION OF ABSTRACT UL		

**Blank**

## PREFACE

The work described in this report was authorized under Project No. 10464806DF97, Sales Order No. 2FK4. This work was started in May 1993 and completed in December 1993.

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

This report has been approved for release to the public. Registered users should request additional copies from the Defense Technical Information Center; unregistered users should direct such requests to the National Technical Information Service.

Accession For	
PTIC ORAL	<input checked="checked" type="checkbox"/>
PTIC WR	<input type="checkbox"/>
UNCLASS	<input type="checkbox"/>
SECRET	<input type="checkbox"/>

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427

1428

1429

1430

1431

1432

1433

1434

1435

1436

1437

1438

1439

1440

1441

1442

1443

1444

1445

1446

1447

1448

1449

1450

1451

1452

1453

1454

1455

1456

1457

1458

1459

1460


14

## QUALITY ASSURANCE

This study, conducted as described by Protocol 22093000X053, was examined for compliance with Good Laboratory Practices as published by the U. S. Environmental Protection Agency in 40 CFR Part 792 (effective 17 Aug 1989). The dates of all inspections and the dates the results of those inspections were reported to the Study Director and management were as follows:

<u>Phase inspected</u>	<u>Date</u>	<u>Date reported</u>
Dosing	20 Jan 93	21 Jan 93
Stock preparation	9 Mar 93	9 Mar 93
Data & Final Report	9 Jun 94	9 Jun 94

To the best of my knowledge, the methods described were the methods followed during the study. The report was determined to be an accurate reflection of the raw data obtained.

  
DENNIS W. JOHNSON  
QA Coordinator, Research & Technology 9 Jun 94

## CONTENTS

	Page
1. INTRODUCTION .....	7
2. METHODS AND MATERIALS .....	8
2.1 Daphnia Assays .....	8
2.2 Fish Assays .....	9
2.3 Earthworm Assay .....	9
2.4 Microtox Assay .....	11
3. RESULTS/DISCUSSION .....	11
4. CONCLUSION .....	13
LITERATURE CITED .....	15

## LIST OF TABLES

1	DS-2 and DS-2P Formulation . . . . .	7
2	Components of the Artificial Soil Used in the Earthworm Toxicity Test . . . . .	10
3	Result from MICROTOX Studies Investigating the Effects of Time on the Toxicity of DS-2 and DS-2P . . . . .	12
4	Toxicity Comparison Between DS-2P and DS-2 . . . . .	14



## AQUATIC TOXICITY OF DECONTAMINATING SOLUTIONS DS-2 / DS-2P

### 1. INTRODUCTION

The Army is working on reducing the possibility of human health effects from exposure to the decontaminating solution DS-2. One of the components of the DS-2 mixture, ethylene glycol monomethyl ether (EGME), has been determined to cause birth defects, fetotoxicity and bone marrow abnormalities in laboratory animals [1]. Eliminating this component will reduce the chance of human health risks and also make transporting and disposal of the solution less restrictive.

The proposed replacement formulation has propylene glycol monomethyl ether (PGME) replacing EGME. Table 1 lists the formulation of DS-2 and the proposed mixture (DS-2P).

Table 1. DS-2 and DS-2P Formulation

<u>DS-2</u>	
Components	Percent by Weight*
Diethylenetriamine	70
Sodium Hydroxide	2
Ethylene glycol monomethyl ether	28
<u>DS-2P</u>	
Diethylenetriamine	70
Sodium Hydroxide	2
Propylene glycol monomethyl ether	28

\* The percent by weight is an average of the manufacturers specification range.

Base line environmental data are needed in order to assess the impact of decon solutions to the environment when used in the field or if accidental spills occur. This study investigates the aquatic toxicity of the DS-2 and DS-2P mixtures on *Daphnia magna*, (water flea), *Pimephales promelas* (fathead minnow), *Eisenia foetida* (earthworm) and *Photobacterium phosphoreum*, (luminescent marine bacterium). The reduction in toxicity of the decon mixtures over time was also investigated using the Microtox Assay (MTX). Toxicity comparisons will determine if replacing EGME with PGME reduces toxic effects of decontamination solutions to the test organisms. The information from this study will support future environmental assessments needed in determining field use rates and disposal management.

Although a number of aquatic organisms are available for short term testing, we chose the daphnia and fathead minnow as the primary test

organisms. These two species are used nationwide, therefore an extensive data base exists for toxicity comparisons. Also, both are inexpensive to culture in the laboratory and can be maintained indefinitely. These factors reduce the variability between organisms and provide the basis for consistently reliable data.

The MTX provides representative species from a different level of biological organization that add to the toxicological predictive power of the screening tests to be performed. The exposure of *P. phosphoreum* (luminescent marine bacterium) to toxicants will typically decrease light output in proportion to increased toxicity, allowing for a dose response relationship to occur. Interest in this assay is based on the following advantages: quick assay time, low cost, small sample size, no organism culturing required, reliability (standardization) and sensitivity. Commercially introduced in 1979, the MTX continues to be studied and evaluated for an array of applications, including toxicity screening of complex effluents, pure compounds, soil sample screening and bioremediation.

One of the first areas impacted during field use and accidental spills is soil, therefore, toxicity studies using the earthworm will yield information on how soil dwelling organisms may be affected from exposure to decon mixtures. The earthworm study determines acute effects (mortality) and subchronic effects (weight loss).

## 2. METHODS AND MATERIALS

All testing conformed to Environmental Protection Agency (EPA) [2,3] and American Society for Testing and Material (ASTM) [4] guidelines. These studies were conducted under Good Laboratory Practice (GLP), and conformed to all inter-agency standard operating procedures.

The decon solutions were received packaged in sealed 1-1/3 quart cans. The cans were opened under a nitrogen atmosphere to prevent CO<sub>2</sub> and moisture contact to the neat solutions. The contents were placed in screw top polycarbonate flasks and sealed with parafilm. All samples were removed from the flasks while under a nitrogen atmosphere. The solutions were tested for reactivity using the chloroform procedures [5] and proved to be within specifications.

### 2.1 Daphnia Assays

The *Daphnia magna* were obtained from Dr. Freida Taub [6] at the University of Washington in Seattle and reared for the past nine years in the laboratory using methods described by Goulden, *et al.* [7]. *Daphnia* stock cultures were fed a mixture of vitamin enriched *Ankistrodesmus falcatus*, *Selenastrum capricornutum* and *Chlamydomonas reinhardtii*. *Daphnia* culture media was derived from well water which was passed through a treatment system containing limestone pH adjustment, iron removal, carbon filtration and UV sterilization. The well water was monitored for 92 commonly found ground water pollutants every six months by Watercheck National Testing Laboratories, Inc.

The test beakers were placed in a temperature controlled room at 20°C with a light-dark cycle of 16:8 hours with 315 ft candles of light. Two replicates per concentration ( 8.0 - 65.0 x 10<sup>-4</sup> % ) containing 10 (*daphnia* less than 24 hours old) in a total of 100 mL of test solution were used. The pH and dissolved

oxygen measurements were taken at the start of testing. After 24 and 48 hours, daphnia were checked for mortality. If the daphnia were not actively swimming, they were touched with a Pasteur pipette. If there were no response or the daphnia could not swim actively for 15 seconds they were considered immobilized. The EC<sub>50</sub> (the effective concentration at which 50 percent of the organisms are immobilized) values were computed using the probit analysis as prepared by Kessler [8]. The EC<sub>50</sub>s were also tabulated graphically using a least square regression analysis verifying all probit results.

## 2.2 Fish Assays

Adult fish were originally obtained from Kurtz's Fish Hatchery in Elverson, PA. These fish were maintained in 40 gallon glass aquaria equipped with under gravel biofilters. The culture water was the same as described in the daphnia methods. Adult fish were fed Tetramin flake food in the mornings and *Lumbriculus variegatus* (black worms) in the afternoon. This feeding regime encouraged the fish to breed continuously. Adult breeding fish are replaced annually to maintain a healthy gene pool.

Adult fish deposit eggs on the under side of clay pots. After the eggs were fertilized, they were transferred to hatchery tanks. Upon hatching, the fry were fed freshly hatched brine shrimp twice a day. The fry were used in toxicity tests after 14 days of age. Fish used in testing were all of similar age and size. The loading did not exceed 0.8 g of fish per liter of solution. If fry appeared stressed or if 5% died within 48 hours before testing, the fry would be discarded. Water temperature was maintained at  $20 \pm 1^\circ\text{C}$  with 16 hours of light and 8 hours of darkness.

The test chambers were one gallon glass jars. The test chambers and glassware were scrubbed with phosphate-free soap, rinsed with tap water until sudsing had ceased, then rinsed with distilled water.

A stock solution of the toxicant was prepared and dispensed directly into the test chambers then diluted to proper concentrations. The dissolved oxygen and pH were measured before fish were transferred to the test chambers. After the fish had been added to the test chambers, a random number table was used to assign each chamber (including controls) to one of two blocks. Next, the chambers were assigned a location number to each of the treatments within that block. (See Appendix 2 for further clarification of randomizing the treatment groups.) The EC<sub>50</sub> values along with the 95% confidence intervals were computed by the probit analysis method and checked graphically using the same procedure described in the daphnia methods.

## 2.3 Earthworm Assay

Earthworm toxicity testing utilized *Eisenia foetida* as the test organism. Survival rates and weight changes were used as indices of toxicity. Test methods used for earthworm toxicity studies were adapted from Karnak and Hamelink [9] and Neuhauser *et al.* [10].

Earthworms were originally purchased from Bert's Bait Farm, Irvine, KY, cultured in a 50/50 mixture of peat moss/potting soil and housed in styrofoam coolers at ambient laboratory temperature.

Earthworms were fed fermented alfalfa pellets obtained by placing the dry alfalfa pellets (commercial rabbit feed) in a sealed container with enough water to cover the pellets. The pellets were periodically mixed and water

added, if necessary, for a two-week period. Ideally, the amount of food added to each cooler should last the earthworms for one week. However, the exact amount added depends on the number and size of the earthworms in each particular cooler. If the food was depleted before one week, additional food was added. The contents of each cooler were mixed once per week to loosen and aerate the soil medium and evenly distribute water throughout the container. Any food remaining on top of the medium was discarded before mixing. After mixing, fresh food was added to the container.

The earthworm toxicity test involved placing 200g substrate and five earthworms into 600-mL glass beaker (two replicated per concentration 500, 1000, 2500, 5000 mg/kg). Earthworm substrate for experiments consisted of a non-sterile artificial soil and distilled water [10]. The heterogeneous parameters of a reproducible artificial soil mixture reduce the variability of the test that could occur if field soil was used. The components of the artificial soil are list in Table 2.

Table 2. Components of the artificial soil used in the Earthworm Toxicity Test.

Stock Components	% by wt.
Lime	1
Finely-ground sphagnum peat moss	10
Kaolinite clay	20
Fine sand	69

The test soil was prepared by mixing (in food blender) the artificial soil with the test substance. Distilled water was slowly added and mixed until a uniform texture was established (25% soil moisture). The test soil was then divided into replicates and placed into 600-mL beakers (200 g of soil).

After the beakers were prepared, 75-100 earthworms were removed from one of the styrofoam coolers and put into a plastic pan. The earthworms were quickly rinsed in tap water and excess water drained from the pan. Five earthworms were randomly picked, quickly blotted with a paper towel, and weighed as a group. They were placed in a beaker which was then covered with nylon screen and cheesecloth secured with a rubber band. The beakers were randomly placed in plastic trays within an incubator. Water was added to the trays to increase the humidity which would reduce drying of the soil in the beakers. The incubator lights were set for continuous operation. Since the earthworms are photophobic, the light encouraged them to burrow into the soil and helped prevent them from crawling out of the beakers.

The earthworms remained in the incubator for a two week exposure period. Beakers were rearranged in the trays at the end of the first week. On day 14, the earthworms were removed from each beaker and reweighed. The earthworms were also examined for changes in color, texture, motility and general physical condition.

Weight change was evaluated using Analysis of Covariance (ANCOVA) to the Newman-Keuls paired comparison of means

## 2.4 Microtox Assay

Materials used in the MTX Assay included lyophilized *Photobacterium phosphoreum* at approximately one hundred million per vial (reagent), 2% sodium chloride solution (diluent) and 22% sodium chloride solution (MTX Osmotic Adjustment Solution, MOAS) for adjustment of osmotic pressure of concentrated samples not requiring pre-dilution with diluent. All materials were supplied by Microbics, Inc.

Use of the Basic Extended Dilution test [11] version of the MTX assay was based on the requirement of an EC<sub>50</sub> endpoint and the ability of the test to encompass a large range of concentrations in which the EC<sub>50</sub> may be found.

The Basic Extended assay was conducted within the temperature controlled (15°C) wells of a photometer (MTX Analyzer). The assay included twelve sample solutions, serially diluted by a factor of two, with three controls. A corresponding set of tubes filled with diluted Reagent (following a 15 minute temperature stabilization period) were read at time zero for initial light output ( $I_0$ ). Aliquots from the corresponding tubes of diluted sample were added and mixed. Light output was then measured at predetermined times (t), usually 5 and 15 minutes. Due to the natural decay of light output over time, the timed readings were normalized using the "Blank Ratio" (BR), which was the ratio of the light output of the control at time t to light output of control at time 0. The BR was applied to  $I_0$ 's to correct for drift and effects of diluting the organisms. The ratio of light lost to light remaining was calculated, and further data reduction produces an EC<sub>50</sub> (the effective concentration at which there is a 50% reduction in light output).

Depending on toxic response over time and quality of data as determined by confidence factors, either the 5 or 15 minute EC<sub>50</sub> was used for comparisons. It is customary to use the 5 minute EC<sub>50</sub> when the values and confidence limits for each time interval are approximately equal. Should data show an increased toxic response for the 15 minute reading and if the 95% confidence range were similar to the 5 minute data, then the 15 minute data was used for comparative purposes. Data for this study is given at t = 5 minutes because all data generated at 15 minutes showed only slight decreases in toxicity, indicating the full effect of the toxicants occurred within 5 minutes.

The DS-2 and DS-2P solutions were diluted separately to obtain the desired stock concentrations in volumetric flasks using 2% sodium chloride (MTX Diluent). All stocks were made within 15 minutes prior to start of assays, with the exception of the assays to determine toxicity reduction over time.

The decon solutions were prepared as previously described and aliquots were taken at time 0, 7 and 14 days for use in determining the toxicity over time. The stocks were maintained under standard laboratory conditions of 8 hours light / 16 hours dark and a temperature of 21°C ± 3° throughout the 14 day analysis. The undiluted material was subjected to the same procedures and the resulting data compared.

## 3. RESULTS/DISCUSSION

Overall, the daphnia were a more sensitive test organism while the earthworms were much less sensitive (having a no effect level of 5000 mg/kg). Substituting PGME for EGME did not significantly change the toxicity of the decon solution.

DS-2 and DS-2P were approximately one order of magnitude less toxic than the DAM decon solution and several orders of magnitude less toxic than malathion (Table 3). The pH of both solutions at 100% was extremely high ( $\text{pH} \geq 13$ , measured using conventional glass electrode). However, the dilutions used during testing lowered the pH to tolerable limits (8.0 - 9.5) [12], thus pH was eliminated as a direct cause of toxicity to daphnia and fish.

The EC<sub>50</sub> values and 95% confidence limits for DS-2P and DS-2 are listed in Table 3. The 95% confidence limits overlap when comparing the toxicity of both solutions.

Stock solutions of DS-2 and DS-2P were prepared by diluting into water and allowed to stand for several days under ambient laboratory conditions. Also, neat samples were allowed to stand under the same conditions. At day 0, 7 and 14, samples were taken and subjected to MICROTOX assays to monitor the change in toxicity. Over a 14 day period, the toxicity of DS-2 and DS-2P did not change (Table 4).

Table 3. Result from MICROTOX studies investigating the effects of time on the toxicity of DS-2 and DS-2P.

	<u>P. phosphoreum</u> 5 min. EC <sub>50</sub> (Vol/Vol %)		
	Day 0	Day 7	Day 14
DS-2P	$2.8 \times 10^{-3}$	$3.5 \times 10^{-3}$	$3.2 \times 10^{-3}$
DS-2	$3.9 \times 10^{-3}$	$4 \times 10^{-3}$	$4.6 \times 10^{-3}$
DAM, [13]	$5.3 \times 10^{-4}$	$5.6 \times 10^{-3}$	$8 \times 10^{-3}$

The specification sheets for the decon solutions require packaging in a moisture and carbon dioxide free atmosphere to prevent degradation. The test results showed no change in toxicity over a 14 day period. This suggest one of three possibilities: the decon solutions did not degrade when exposed to moisture, the degradation occurred instantly and the toxicity of non-degraded decon solution could never be obtained in an aquatic environment, or the materials were degraded before opening due to improper packaging.

The activity of the decon solutions met specs after being divided (under nitrogen) into smaller samples, therefore eliminating the possibility of improper packaging.

When diluted to approximately 50 % with water, DS-2 degrades instantaneously [14]. Assuming DS-2P reacts similarly to DS-2 when added to water, complete degradation may have occurred before toxicity end point were reached.

The most probable reason for not seeing a change in toxicity over time is the instantaneous degradation of the decon solution when added to water. Even though the MICROTOX end points are reached in 5 minutes, a change in toxicity was not observed.

It is assumed the degradation of DS-2P is instantaneous (similar to DS-2 in water) and the toxicity estimates presented in this paper are associated with the by-products produced when adding the decon solutions to water.

#### 4. CONCLUSION

The substitution of PGME for EGME did not significantly change the toxicity of the decon solutions to the test organisms.

Over time, the toxicity of DS-2P did not change. Therefore it is assumed that the toxicity estimates of the decon solutions are directly related to the degradation by-products produced when added to water. Fate studies are needed to confirm the reaction that occurs when adding DS-2P to water.

The decon solutions scored a ranking of 9, highly toxic (out of a scale of 0 - 9, 9 being the most toxic), on the Chemical scoring System for Hazard and Exposure Identification [17].

If used in the open environment, efforts should be made to provide as much containment of decon solutions as possible to prevent runoff into the surrounding ecosystem.

Table 4. Toxicity Comparison Between DS-2P and DS-2

	Daphnia 48 hr EC50 (95% CI) Vol/Vol %	Fathead Minnow 96 hr EC50 (95% CI) Vol/Vol %	P. phosphoreum 5 min EC50 (95% CI) Vol/Vol %	Earthworm 14 day EC50 (95% CI) mg/kg
DS-2P	$1.4 \times 10^{-3}$ (1.2 - 1.7)	$2.8 \times 10^{-2}$ (2.0 - 3.0)	$4 \times 10^{-3}$ (2 - 9)	N.E. *
DS-2	$1.7 \times 10^{-3}$ (1.3 - 2.1)	$2.2 \times 10^{-2}$ (2.0 - 3.0)	$3 \times 10^{-3}$ (2 - 4)	N.E.
D.A.M. [14]	$5.0 \times 10^{-5}$ (4.8 - 5.2)	-----	$5 \times 10^{-4}$ (4 - 7)	-----
Malathion	$8.0 \times 10^{-8}$ , [15]	$8.0 \times 10^{-4}$ , [15]	$2.4 \times 10^{-6}$ , [16]	

\* N.E. - No effects were seen up to 2,500 mg/kg.



# LITERATURE CITED

1. Sigma Aldrich Corporation, Material Safety Data Sheet, 1001 West Saint Paul Ave., Milwaukee, WI 53233, 1989.
2. U.S. Environmental Protection Agency, Users guide: procedures for conducting *Daphnia magna* toxicity bioassays. EPA-660/8-87/011. March 1987
3. U.S. Environmental Protection Agency, Methods for acute toxicity tests with fish, macroinvertebrates and amphibians, EPA 660/3-75-009, 1975.
4. American Society for Testing and Materials. Standard E729, Guide for conducting acute toxicity tests with fishes, macroinvertebrates and amphibians, 1986.
5. Analysis provided by Vincent Auber CBDCOM, using procedures listed in the Military Specification for Decontaminating Agent, DS2, MIL-D-50030G(EA), 2 June 1986.
6. Taub, F.B. and Read, P.L., Standardized aquatic microcosm protocol, Vol. II, Final Report Contract No. 223-80-2352, Food and Drug Administration, Washington, DC, 1983.
7. Goulden, C.E., R.M. Conotto, J.A. Hendrickson, Jr., L.L. Hornig and K.L. Johnson, Procedures and recommendations for the culture and use of *Daphnia magna* bioassay studies. ASTM Special Technical Report Publication 766, p 139-160, 1982
8. Kessler, F., Probit analysis, U.S. EPA, 26 West Martin Luther King Dr., Cincinnati, Ohio 45268.
9. Karnak, R. and J. Hamelink, A standardized method for determining acute toxicity of chemicals to earthworms, *Ecotox. Environ. Safety*, Vol 6, p 216-222, 1982.
10. Neuhauser, E. F., R. C. Lochr, M. R. Malecki, D. L. Milligan and P. R. Durkin, The toxicity of selected organic chemicals to the earthworm *Eisenia fetida*, *J. Environmental Quality*, Vol 14, p 383-388, 1985.
12. Chester, N. A., Haley, M. V., and Landis, W. G., The aquatic toxicity of isopropylamine: comparison of experimentally derived values with structure activity predictions, *Environmental Sciences*, 1, 3, p 117-126, 1982.
13. Haley, M. V., Kurnas, C. W., Chester, N. A. and Muse, W. T., The aquatic toxicity of the decontaminating agent: multipurpose (DAM) decontamination solution, Technical Report CBDA-TR-149, June 1993.
14. Beaudry, W. T., Szafraniec, L. L., and Leslie, D. R., Reactions of chemical warfare agents with DS-2: Product identification by NMR organophosphorous compounds, CRDEC-TR-364, June, 199

15. Johnson, W.W. and Finley, M.T., Hand Book of Acute Toxicity of Chemicals to Fish and Aquatic Invertebrates. United States Department of the Interior Fish and Wildlife Service, Resource Publication 137, Washington, D.C., 1980.
16. Kaiser, K. L. E., Photobacterium phosphoreum toxicity bioassay, II. toxicity data compilation, Toxicity Assessment: An International Journal, Vol 3, p 195-227, 1988.
17. O'Bryan, T. R. and R. H. Ross, Chemical scoring system for hazard and exposure identification, J. of Toxicology and Environmental Health, Vol 1, p 119-134, 1988.

**SUPPLEMENTARY**

**INFORMATION**

DEPARTMENT OF THE ARMY  
U.S. Army Edgewood Research, Development and Engineering Center  
Aberdeen Proving Ground, Maryland 21010-5423

ERRATUM SHEET

8 November 1994

Report No. ERDEC-TR-202  
Title Aquatic Toxicity of Decontaminating Solutions  
DS-2/DS-2P  
Authors M.V. Haley, N.A. Chester, C.W. Kurnas, and C.T. Phillips  
Date September 1994  
Classification UNCLASSIFIED

In the document recently distributed to you, please replace pages 9, 10, 15, and 16 with the attached corrected pages.

1 Atch  
as stated

*Sandra J. Johnson*

SANDRA J. JOHNSON  
Chief, Technical Releases Office

ERRATA ADA-2859920

oxygen measurements were taken at the start of testing. After 24 and 48 hours, daphnia were checked for mortality. If the daphnia were not actively swimming, they were touched with a Pasteur pipette. If there were no response or the daphnia could not swim actively for 15 seconds they were considered immobilized. The EC<sub>50</sub> (the effective concentration at which 50 percent of the organisms are immobilized) values were computed using the probit analysis as prepared by Kessler [8]. The EC<sub>50</sub>s were also tabulated graphically using a least square regression analysis verifying all probit results.

## 2.2 Fish Assays

Adult fish were originally obtained from Kurtz's Fish Hatchery in Elverson, PA. These fish were maintained in 40 gallon glass aquaria equipped with under gravel biofilters. The culture water was the same as described in the daphnia methods. Adult fish were fed Tetramin flake food in the mornings and *Lumbriculus variegatus* (black worms) in the afternoon. This feeding regime encouraged the fish to breed continuously. Adult breeding fish are replaced annually to maintain a healthy gene pool.

Adult fish deposit eggs on the under side of clay pots. After the eggs were fertilized, they were transferred to hatchery tanks. Upon hatching, the fry were fed freshly hatched brine shrimp twice a day. The fry were used in toxicity tests after 14 days of age. Fish used in testing were all of similar age and size. The loading did not exceed 0.8 g of fish per liter of solution. If fry appeared stressed or if 5% died within 48 hours before testing, the fry would be discarded. Water temperature was maintained at  $20 \pm 1^\circ\text{C}$  with 16 hours of light and 8 hours of darkness.

The test chambers were one gallon glass jars. The test chambers and glassware were scrubbed with phosphate-free soap, rinsed with tap water until sudsing had ceased, then rinsed with distilled water.

A stock solution of the toxicant was prepared and dispensed directly into the test chambers then diluted to proper concentrations. The dissolved oxygen and pH were measured before fish were transferred to the test chambers. After the fish had been added to the test chambers, a random number table was used to assign each chamber (including controls) to one of two blocks. Next, the chambers were assigned a location number to each of the treatments within that block. The EC<sub>50</sub> values along with the 95% confidence intervals were computed by the probit analysis method and checked graphically using the same procedure described in the daphnia methods.

## 2.3 Earthworm Assay

Earthworm toxicity testing utilized *Eisenia foetida* as the test organism. Survival rates and weight changes were used as indices of toxicity. Test methods used for earthworm toxicity studies were adapted from Karnak and Hamelink [9] and Neuhauser *et al.* [10].

Earthworms were originally purchased from Bert's Bait Farm, Irvine, KY, cultured in a 50/50 mixture of peat moss/potting soil and housed in styrofoam coolers at ambient laboratory temperature.

Earthworms were fed fermented alfalfa pellets obtained by placing the dry alfalfa pellets (commercial rabbit feed) in a sealed container with enough water to cover the pellets. The pellets were periodically mixed and water

added, if necessary, for a two-week period. Ideally, the amount of food added to each cooler should last the earthworms for one week. However, the exact amount added depends on the number and size of the earthworms in each particular cooler. If the food was depleted before one week, additional food was added. The contents of each cooler were mixed once per week to loosen and aerate the soil medium and evenly distribute water throughout the container. Any food remaining on top of the medium was discarded before mixing. After mixing, fresh food was added to the container.

The earthworm toxicity test involved placing 200g substrate and five earthworms into 600-mL glass beaker (two replicated per concentration 500, 1000, 2500, 5000 mg/kg). Earthworm substrate for experiments consisted of a non-sterile artificial soil and distilled water [10]. The heterogeneous parameters of a reproducible artificial soil mixture reduce the variability of the test that could occur if field soil was used. The components of the artificial soil are list in Table 2.

Table 2. Components of the artificial soil used in the Earthworm Toxicity Test.

Stock Components	% by wt.
Lime	1
Finely-ground sphagnum peat moss	10
Kaolinite clay	20
Fine sand	69

The test soil was prepared by mixing (in food blender) the artificial soil with the test substance. Distilled water was slowly added and mixed until a uniform texture was established (25% soil moisture). The test soil was then divided into replicates and placed into 600-mL beakers (200 g of soil).

After the beakers were prepared, 75-100 earthworms were removed from one of the styrofoam coolers and put into a plastic pan. The earthworms were quickly rinsed in tap water and excess water drained from the pan. Five earthworms were randomly picked, quickly blotted with a paper towel, and weighed as a group. They were placed in a beaker which was then covered with nylon screen and cheesecloth secured with a rubber band. The beakers were randomly placed in plastic trays within an incubator. Water was added to the trays to increase the humidity which would reduce drying of the soil in the beakers. The incubator lights were set for continuous operation. Since the earthworms are photophobic, the light encouraged them to burrow into the soil and helped prevent them from crawling out of the beakers.

The earthworms remained in the incubator for a two week exposure period. Beakers were rearranged in the trays at the end of the first week. On day 14, the earthworms were removed from each beaker and reweighed. The earthworms were also examined for changes in color, texture, motility and general physical condition.

Weight change was evaluated using Analysis of Covariance (ANCOVA) to the Newman-Keuls paired comparison of means.

#### LITERATURE CITED

1. Sigma Aldrich Corporation, Material Safety Data Sheet, 1001 West Saint Paul Ave., Milwaukee, WI 53233, 1989.
2. U.S. Environmental Protection Agency, Users guide: procedures for conducting *Daphnia magna* toxicity bioassays. EPA-660/8-87/011. March 1987
3. U.S. Environmental Protection Agency, Methods for acute toxicity tests with fish, macroinvertebrates and amphibians, EPA 660/3-75-009, 1975.
4. American Society for Testing and Materials. Standard E729, Guide for conducting acute toxicity tests with fishes, macroinvertebrates and amphibians, 1986.
5. Analysis provided by Vincent Auber CBDCOM, using procedures listed in the Military Specification for Decontaminating Agent, DS2, MIL-D-50030G(EA), 2 June 1986.
6. Taub, F.B. and Read, P.L., Standardized aquatic microcosm protocol, Vol. II, Final Report Contract No. 223-80-2352, Food and Drug Administration, Washington, DC, 1983.
7. Goulden, C.E., R.M. Conotto, J.A. Hendrickson, Jr., L.L. Hornig and K.L. Johnson, Procedures and recommendations for the culture and use of *Daphnia magna* in bioassay studies. ASTM Special Technical Report Publication 766, p 139-160, 1982
8. Kessler, F., Probit analysis, U.S. EPA, 26 West Martin Luther King Dr., Cincinnati, Ohio 45268.
9. Karnak, R. and J. Hamelink, A standardized method for determining acute toxicity of chemicals to earthworms, *Ecotox. Environ. Safety*, Vol 6, p 216-222, 1982.
10. Neuhauser, E. F., R. C. Loehr, M. R. Malecki, D. L. Milligan and P. R. Durkin, The toxicity of selected organic chemicals to the earthworm *Eisenia fetida*, *J. Environmental Quality*, Vol 14, p 383-388, 1985.
11. MICROTOX Model 500 Toxicity Test System Manual, Microbics Corporation, 2232 Rutherford Rd., Carlsbad, CA 92008, P 37, 5-30-91, May 1991
12. Chester, N. A., Haley, M. V., and Landis, W. G., The aquatic toxicity of isopropylamine: comparison of experimentally derived values with structure activity predictions, *Environmental Sciences*, 1, 3, p 117-126, 1992.
13. Haley, M. V., Kurnas, C. W., Chester, N. A. and Muse, W. T., The aquatic toxicity of the decontaminating agent: multipurpose (DAM) decontamination solution, Technical Report CBDA-TR-149, June 1993.

14. Beaudry, W. T., Szafraniec, L. L., and Leslie, D. R., Reactions of chemical warfare agents with DS-2: Product identification by NMR organophosphorous compounds, CRDEC-TR-364, June, 199
15. Johnson, W.W. and Finley, M.T., Hand Book of Acute Toxicity of Chemicals to Fish and Aquatic Invertebrates, United States Department of the Interior Fish and Wildlife Service, Resource Publication 137, Washington, D.C., 1980.
16. Kaiser, K. L. E., Photobacterium phosphoreum toxicity bioassay, II. toxicity data compilation, Toxicity Assessment: An International Journal, Vol 3, p 195-237, 1988.
17. O'Bryan, T. R. and R. H. Ross, Chemical scoring system for hazard and exposure identification, J. of Toxicology and Environmental Health, Vol 1, p 119-134, 1988.